

Explicit and Exact Solutions to a Kolmogorov-Petrovskii-Piskunov Equation

W. X. Ma¹ and B. Fuchssteiner

FB17, Mathematik-Informatik, Universitaet Paderborn, D-33098 Paderborn, Germany

Abstract

Some explicit traveling wave solutions to a Kolmogorov-Petrovskii-Piskunov equation are presented through two ansätze. By a Cole-Hopf transformation, this Kolmogorov-Petrovskii-Piskunov equation is also written as a bilinear equation and further two solutions to describe nonlinear interaction of traveling waves are generated. Bäcklund transformations of the linear form and some special cases are considered.

1 Introduction

The Cauchy problem of the Kolmogorov-Petrovskii-Piskunov equation [13]

$$\begin{cases} u_t - u_{xx} = f(u), & f \text{ nonlinear, } f(0) = 0, \\ u(x, 0) = \phi(x), & x \in R^1 \end{cases}$$

has been extensively investigated both by analytic techniques [3], [10], and by probabilistic methods [18], [4], and the existence of traveling wave solutions with various velocities has been also proved. A special case is the Fisher equation

$$u_t - u_{xx} = u(1 - u),$$

which was originally proposed [7] as a model for the propagation of a favored gene. An explicit and exact solitary solution of the Fisher equation may be presented by Painlevé analysis [2]. As an example of the above Kolmogorov-Petrovskii-Piskunov equation, Kametaka [10] considered the Cauchy problem of a generalized Fisher equation

$$\begin{cases} u_t - u_{xx} = \lambda_0^2 u(1 - u^n), \\ u(x, 0) = \{1 + (2^{n/2} - 1)e^{-(n/2)\sigma x}\}^{-2/n}, & x \in R^1, \end{cases}$$

where $\lambda_0 > 0$, $n \in N$, $\sigma > 0$, and gave an explicit solution

$$u(x, t) = \left[1 + (2^{n/2} - 1)\exp\left\{-\frac{n}{2}\sigma_1(x + 2\lambda_1 t)\right\} \right]^{-2/n}$$

when

$$\sigma = \sigma_1 = \lambda_1 - \sqrt{\lambda_1^2 - \lambda_0^2}, \quad \lambda_1 = \frac{1}{2} \left\{ \left(\frac{n}{2} + 1\right)^{1/2} + \left(\frac{n}{2} + 1\right)^{-1/2} \right\} \lambda_0.$$

Abdelkader [1] and Wang [21] extended the integer n to a real number α satisfying $\alpha \geq 1$ and Wang obtained a class of explicit traveling wave solutions successfully by introducing a special nonlinear transformation.

¹On leave of absence from Institute of Mathematics, Fudan University, Shanghai 200433, China

In this paper, we consider the following Kolmogorov-Petrovskii-Piskunov equation

$$u_t - u_{xx} + \mu u + \nu u^2 + \delta u^3 = 0 \quad (1)$$

where μ, ν, δ are three real constants. Some special cases with $\mu + \nu + \delta = 0$ of this equation have been studied (see for instance [6], [12], [9] and [14]). Our purpose is to look for explicit and exact solutions for the general case of (1). Note that a more general equation

$$u_{t'} - \alpha u_{x'x'} + \mu u + \nu u^2 + \delta u^3 = 0, \quad \alpha \neq 0$$

may be mapped into the Kolmogorov-Petrovskii-Piskunov equation (1) by a time scaling $t' \rightarrow \alpha t$ and therefore the Kolmogorov-Petrovskii-Piskunov equation (1) is no loss of generality. In Section 2, we analyze the possibilities of solutions which correspond to two Riccati equations and further give explicitly a number of exact solutions to (1). The essential point is to break the nonlinear equation (1) into two smaller problems and then to solve these two smaller problems. In Section 3, we change the Kolmogorov-Petrovskii-Piskunov (1) into a bilinear equation, like the Hirota bilinear one, by making the well known Cole-Hopf transformation, and present two explicit solutions to describe nonlinear coalescence of traveling wave solutions. In Section 4, Bäcklund transformations of the linear form are discussed along with some explicit relations of Bäcklund transformations for the obtained solutions. In Section 5, more discussion is given.

2 Traveling wave solutions

We consider traveling wave solutions to the Kolmogorov-Petrovskii-Piskunov (1)

$$u(x, t) = u(\xi) = u(kx - \omega t),$$

where the wavenumber k and the frequency ω are required to be nonzero for generating nontrivial solutions. The resulting ordinary differential equation from (1) reads as

$$-\omega u_\xi - k^2 u_{\xi\xi} + \mu u + \nu u^2 + \delta u^3 = 0. \quad (2)$$

In the following we generate traveling wave solutions to (1) starting with two ansätze.

First we make the ansatz for (2)

$$u(\xi) = \sum_{i=0}^M a_i v^i = \sum_{i=0}^M a_i (v(\xi))^i, \quad v_\xi = \varepsilon(1 - v^2), \quad \varepsilon = \pm 1. \quad (3)$$

It is easy to show that M must be 1 if the functions $1, v, v^2, \dots, v^m$ ($m \in \mathbb{N}$) are linear independent and $\delta \neq 0$. So without loss of generality, we may choose

$$u = a_0 + a_1 v,$$

and thus

$$\begin{aligned} u_\xi &= \varepsilon a_1 - \varepsilon a_1 v^2, \\ u_{\xi\xi} &= -2a_1 v + 2a_1 v^3. \end{aligned}$$

The substitution into (2) yields the following conditions for determining a_0, a_1, k, ω :

$$\begin{cases} -\varepsilon \omega a_1 + \mu a_0 + \nu a_0^2 + \delta a_0^3 = 0, \\ 2k^2 a_1 + \mu a_1 + 2\nu a_0 a_1 + 3\delta a_0^2 a_1 = 0, \\ \varepsilon \omega a_1 + \nu a_1^2 + 3\delta a_0 a_1^2 = 0, \\ -2k^2 a_1 + \delta a_1^3 = 0. \end{cases} \quad (4)$$

We need to assume $a_1 \neq 0$ for nontrivial solutions, and thereby we get

$$k^2 = \frac{1}{2}\delta a_1^2, \quad \omega = -\varepsilon(\nu a_1 + 3\delta a_0 a_1), \quad (5)$$

$$a_1^2 = -\frac{\mu}{\delta} - \frac{2\nu}{\delta}a_0 - 3a_0^2, \quad (6)$$

$$f(a_0) := \frac{\mu\nu}{\delta} + 2\left(\frac{\nu^2}{\delta} + \mu\right)a_0 + 8\nu a_0^2 + 8\delta a_0^3 = 0, \quad (7)$$

in which (5) shows that $\delta > 0$. The equation (7) is a cubic equation. Krishnan [14] analyzed the case of (7) with $\mu = a$, $\nu = -(a + 1)$, $\delta = 1$, but he failed to give any solution. Actually, this equation has a solution $a_0 = -\frac{\nu}{2\delta}$ by inspection. We can accordingly decompose

$$f(a_0) = \left(a_0 + \frac{\nu}{2\delta}\right)(8\delta a_0^2 + 4\nu a_0 + 2\mu),$$

from which we acquire three roots

$$a_{01} = -\frac{\nu}{2\delta}, \quad a_{02,3} = \frac{-\nu \pm \sqrt{\Delta}}{4\delta}, \quad \Delta = \nu^2 - 4\mu\delta. \quad (8)$$

Of course, we can also solve the cubic equation (7) by computer algebra tools, for example, Mathematica, Maple and MuPAD. Further we can obtain by (6) and (5)

$$a_{11} = \varepsilon_1 \frac{\sqrt{\Delta}}{2\delta}, \quad a_{12,3} = \varepsilon_{2,3} a_{02,3}; \quad (9)$$

$$k_1 = \varepsilon_4 \frac{\sqrt{\Delta}}{2\sqrt{2\delta}}, \quad k_2 = \varepsilon_5 \frac{\nu - \sqrt{\Delta}}{4\sqrt{2\delta}}, \quad k_3 = \varepsilon_6 \frac{\nu + \sqrt{\Delta}}{4\sqrt{2\delta}}; \quad (10)$$

$$\omega = \varepsilon\varepsilon_1 \frac{\nu\sqrt{\Delta}}{4\delta}, \quad \omega_{2,3} = -\varepsilon\varepsilon_{2,3} \left[\frac{(\nu \mp \sqrt{\Delta})^2}{16\delta} - \frac{\nu}{2} \right]. \quad (11)$$

Now we may conclude that only when $\delta > 0$, $\Delta \geq 0$, there are real solutions a_0, a_1, k, ω and we acquire the following three exact solutions for the Kolmogorov-Petrovskii-Piskunov equation (1) with $\delta > 0$, $\Delta \geq 0$:

$$\begin{aligned} u_1(x, t) &= u_1(x, t; \varepsilon_1, \varepsilon_4) = a_{01} + a_{11}v(k_1x - \omega_1t) \\ &= -\frac{\nu}{2\delta} + \varepsilon_1 \frac{\sqrt{\Delta}}{2\delta} v\left(\varepsilon_4 \frac{\sqrt{\Delta}}{2\sqrt{2\delta}}x - \varepsilon_1 \frac{\nu\sqrt{\Delta}}{4\delta}t\right), \end{aligned} \quad (12)$$

$$\begin{aligned} u_2(x, t) &= u_2(x, t; \varepsilon_2, \varepsilon_5) = a_{02} + a_{12}v(k_2x - \omega_2t) \\ &= \frac{-\nu + \sqrt{\Delta}}{4\delta} + \varepsilon_2 \frac{-\nu + \sqrt{\Delta}}{4\delta} v\left(\varepsilon_5 \frac{\nu - \sqrt{\Delta}}{4\sqrt{2\delta}}x + \varepsilon_2 \left[\frac{(\nu - \sqrt{\Delta})^2}{16\delta} - \frac{\mu}{2}\right]t\right), \end{aligned} \quad (13)$$

$$\begin{aligned} u_3(x, t) &= u_3(x, t; \varepsilon_3, \varepsilon_6) = a_{03} + a_{13}v(k_3x - \omega_3t) \\ &= \frac{-\nu - \sqrt{\Delta}}{4\delta} + \varepsilon_3 \frac{-\nu - \sqrt{\Delta}}{4\delta} v\left(\varepsilon_6 \frac{\nu + \sqrt{\Delta}}{4\sqrt{2\delta}}x + \varepsilon_3 \left[\frac{(\nu + \sqrt{\Delta})^2}{16\delta} - \frac{\mu}{2}\right]t\right), \end{aligned} \quad (14)$$

where $v_\xi = 1 - v^2$, $\varepsilon_i = \pm 1$, $1 \leq i \leq 6$. In the above solutions, we cancel the case of $\varepsilon = -1$ due to the same solutions.

Notice that the Riccati equation $v_\xi = a(1 - v^2)$ ($a \in R^1$) has a general solution

$$v = v(\xi) = \frac{A - Be^{-2a\xi}}{A + Be^{-2a\xi}} = \begin{cases} 1 & \text{for } B = 0, \\ -1 & \text{for } A = 0, \\ \tanh\left(a\xi - \frac{1}{2}\ln\left(\frac{B}{A}\right)\right) & \text{for } AB > 0, \\ \coth\left(a\xi - \frac{1}{2}\ln\left(-\frac{B}{A}\right)\right) & \text{for } AB < 0, \end{cases} \quad (15)$$

where A, B are arbitrary constants satisfying $A^2 + B^2 \neq 0$. This solution may be obtained by three tricks: a Möbius transformation, a Cole-Hopf transformation or a relation [11]

$$\frac{(v_1 - v_2)(v_3 - v_4)}{(v_1 - v_3)(v_2 - v_4)} = C, \quad C = \text{const.}$$

of the solutions v_i , $1 \leq i \leq 4$, beginning with three known solutions $1, -1, \tanh(\xi)$. From $v(\xi) = \pm 1$, (12), (13) and (14) result in three constant solutions of (1), but after choosing

$$v(\xi) = \tanh(\xi + \xi_0), \quad v(\xi) = \coth(\xi + \xi_0) \quad (\xi_0 \text{ arbitrary}),$$

(12), (13) and (14) yield non trivial solutions: three explicit traveling front solutions and three explicit singular traveling solutions, respectively.

Secondly we make another ansatz for (2)

$$u(\xi) = \sum_{i=0}^M b_i v^i = \sum_{i=0}^M b_i (v(\xi))^i, \quad v_\xi = \varepsilon(1 + v^2), \quad \varepsilon = \pm 1. \quad (16)$$

Similarly M must equal 1 if the functions $1, v, v^2, \dots, v^m$ ($m \in N$) and $\delta \neq 0$. So without loss of generality, we can choose

$$u = b_0 + b_1 v,$$

and further we find

$$\begin{aligned} u_\xi &= \varepsilon b_1 + \varepsilon b_1 v^2, \\ u_{\xi\xi} &= 2b_1 v + 2b_1 v^3. \end{aligned}$$

The substitution into (2) engenders the following conditions on b_0, b_1, k, ω :

$$\begin{cases} -\varepsilon\omega b_1 + \mu b_0 + \nu b_0^2 + \delta b_0^3 = 0, \\ -2k^2 b_1 + \mu b_1 + 2\nu b_0 b_1 + 3\delta b_0^2 b_1 = 0, \\ -\varepsilon\omega b_1 + \nu b_1^2 + 3\delta b_0 b_1^2 = 0, \\ -2k^2 b_1 + \delta b_1^3 = 0. \end{cases} \quad (17)$$

Note that there are only two terms in (4) and (17) have opposite signs. In an analogous way, we can prove that only when $\delta > 0$, $\Delta \leq 0$, there exists a set of real nonzero solutions b_0, b_1, k, ω , which may be worked out

$$b_0 = -\frac{\nu}{2\delta}, \quad b_1 = \varepsilon_1 \frac{\sqrt{-\Delta}}{2\delta}, \quad k = \varepsilon_2 \frac{\sqrt{-\Delta}}{2\sqrt{2\delta}}, \quad \omega = -\varepsilon\varepsilon_1 \frac{\nu\sqrt{-\Delta}}{4\delta}. \quad (18)$$

In this case, notice that the corresponding Riccati equation $v_\xi = a(1 + v^2)$, $a \in R^1$ has the solutions

$$v(\xi) = \tan(a\xi + \xi_0), \quad v(\xi) = -\cot(a\xi + \xi_0)$$

with an arbitrary constant ξ_0 . Accordingly we obtain two explicit exact solutions for the Kolmogorov-Petrovskii-Piskunov (1) with $\delta > 0$, $\Delta \leq 0$:

$$u'_1 = -\frac{\nu}{2\delta} + \varepsilon_1 \frac{\sqrt{-\Delta}}{2\delta} \tan\left(\varepsilon_2 \frac{\sqrt{-\Delta}}{2\sqrt{2\delta}} x + \varepsilon_1 \frac{\nu\sqrt{-\Delta}}{4\delta} t + \xi_0\right), \quad (19)$$

$$u'_2 = -\frac{\nu}{2\delta} - \varepsilon_1 \frac{\sqrt{-\Delta}}{2\delta} \cot\left(\varepsilon_2 \frac{\sqrt{-\Delta}}{2\sqrt{2\delta}} x + \varepsilon_1 \frac{\nu\sqrt{-\Delta}}{4\delta} t + \xi_0\right), \quad (20)$$

where $\varepsilon_1, \varepsilon_2 = \pm 1$ and ξ_0 is arbitrary. In the above solutions, ε is again incorporated into ε_1 due to the same solution.

3 Nonlinear interactions of traveling waves

We make a Cole-Hopf transformation

$$u = \alpha(\ln f)_x = \alpha f_x/f, \quad \alpha = \text{const.} \quad (21)$$

for the Kolmogorov-Petrovskii-Piskunov equation (1). We have

$$(f_{xt}f - f_x f_t)f^2 - (f_{xxx}f - 3f_x f_{xx})f^2 - 2f_x^3 f + \mu f_x f^3 + \nu \alpha f_x^2 f^2 + \delta \alpha^2 f_x^3 f = 0. \quad (22)$$

Therefore if we choose $\alpha = \pm\sqrt{2/\delta} = \varepsilon\sqrt{2/\delta}$, we get a bilinear equation

$$f_{xt}f - f_x f_t - f_{xxx}f + 3f_x f_{xx} + \mu f_x f + \nu \alpha f_x^2 = 0. \quad (23)$$

After assuming a kind of solutions to be expressed by an exponential function

$$f = A_1 e^{k_1 x + \omega_1 t} + A_2 e^{k_2 x + \omega_2 t} + A_3 e^{k_3 x + \omega_3 t},$$

we find the conditions

$$\begin{cases} 2k_i^3 + \nu \alpha k_i^2 + \mu k_i = 0, & 1 \leq i \leq 3, \\ (\omega_i - \omega_j)(k_i - k_j) - (k_i^3 + k_j^3 - 3k_i k_j^2 - 3k_i^2 k_j) \\ \quad + \mu(k_i + k_j) + 2\nu \alpha k_i k_j = 0, & 1 \leq i < j \leq 3. \end{cases} \quad (24)$$

By solving this equation, we get a nontrivial solution to the equation (23)

$$f = A_1 + A_2 e^{k_+ x + (k_+^2 - \mu)t} + A_3 e^{k_- x + (k_-^2 - \mu)t}, \quad (25)$$

where A_i , $1 \leq i \leq 3$, are arbitrary constants and

$$k_{\pm} = \frac{-\varepsilon\nu \pm \sqrt{\Delta}}{2\sqrt{2\delta}} = \frac{-\varepsilon\nu \pm \sqrt{\nu^2 - 4\mu\delta}}{2\sqrt{2\delta}}.$$

Finally we obtain two different solutions to the Kolmogorov-Petrovskii-Piskunov equation (1) with $\delta > 0$, $\Delta \geq 0$:

$$u_4(x, t) = u_4(x, t; A_1, A_2, A_3) = \sqrt{\frac{2}{\delta}} \frac{A_2 k_+ e^{\eta_+} + A_3 k_- e^{\eta_-}}{A_1 + A_2 e^{\eta_+} + A_3 e^{\eta_-}}, \quad (26)$$

with

$$\begin{aligned} k_{\pm} &= \frac{-\nu \pm \sqrt{\Delta}}{2\sqrt{2\delta}}, \quad \eta_{\pm} = k_{\pm} x + (k_{\pm}^2 - \mu)t; \\ u_5(x, t) &= u_5(x, t; A_1, A_2, A_3) = -\sqrt{\frac{2}{\delta}} \frac{A_2 k_+ e^{\eta_+} + A_3 k_- e^{\eta_-}}{A_1 + A_2 e^{\eta_+} + A_3 e^{\eta_-}}, \end{aligned} \quad (27)$$

with

$$k_{\pm} = \frac{\nu \pm \sqrt{\Delta}}{2\sqrt{2\delta}}, \quad \eta_{\pm} = k_{\pm} x + (k_{\pm}^2 - \mu)t;$$

where A_1, A_2, A_3 are three arbitrary constants. We note that $u_4(-x, t, A_1, A_2, A_3) = u_5(x, t, A_1, A_3, A_2)$. Therefore for the case of FitzHugh-Nagumo equation, u_5 is exactly a solution lost in Ref. [12]. The solutions u_4, u_5 describe the coalescence of two traveling fronts or two singular traveling waves of the same sense. Direct numerical calculations of nonlinear interactions for the FitzHugh-Nagumo case of (26) were done in Ref. [12]. These two solutions are analytic on the whole plane of (x, t) when $A_i A_j > 0$, $i \neq j$, and they blow up at some points of (x, t) when A_i , $1 \leq i \leq 3$, don't possess the same signs.

4 Bäcklund transformations

We know there are three solutions

$$\beta = 0, \beta = C_{\pm} = \frac{-\nu \pm \sqrt{\Delta}}{2\delta}$$

to the equation $\mu\beta + \nu\beta^2 + \delta\beta^3 = 0$, when $\Delta = \nu^2 - 4\mu\delta \geq 0$. Make a linear transformation

$$u = \alpha\tilde{u} + \beta, \alpha \neq 0, \beta = 0 \text{ or } C_{\pm}. \quad (28)$$

This moment

$$\begin{aligned} \mu u + \nu u^2 + \delta u^3 &= \alpha(\mu + 2\beta\nu + 3\beta^2\delta)\tilde{u} + \alpha^2(\nu + 3\beta\delta)\tilde{u}^2 + \alpha^3\delta\tilde{u}^3 \\ &= \begin{cases} \alpha\mu\tilde{u} + \alpha^2\nu\tilde{u}^2 + \alpha^3\delta\tilde{u}^3, & \text{for } \beta = 0, \\ -\alpha(2\mu + \beta\nu)\tilde{u} + \alpha^2(\nu + 3\beta\delta)\tilde{u}^2 + \alpha^3\delta\tilde{u}^3, & \text{for } \beta = C_{\pm}. \end{cases} \end{aligned}$$

and thus the Kolmogorov-Petrovskii-Piskunov equation (1) is equivalent to the following new Kolmogorov-Petrovskii-Piskunov equation

$$\begin{aligned} \tilde{u}_t - \tilde{u}_{xx} &= -(\mu + 2\beta\nu + 3\beta^2\delta)\tilde{u} - \alpha(\nu + 3\beta\delta)\tilde{u}^2 - \alpha^2\delta\tilde{u}^3 \\ &:= -\tilde{\mu}\tilde{u} - \tilde{\nu}\tilde{u}^2 - \tilde{\delta}\tilde{u}^3. \end{aligned} \quad (29)$$

A direct calculation yields that

$$\tilde{\Delta} := \tilde{\nu}^2 - \tilde{\mu}\tilde{\delta} = \begin{cases} \alpha^2\Delta & \text{when } \beta = 0; \\ \frac{1}{4}\alpha^2(\nu + \sqrt{\Delta})^2, & \text{when } \beta = C_+; \\ \frac{1}{4}\alpha^2(\nu - \sqrt{\Delta})^2, & \text{when } \beta = C_-. \end{cases} \quad (30)$$

We remark that a similar equation $\tilde{u}_t - \tilde{u}_{xx} + \mu\tilde{u} - \nu\tilde{u}^2 + \delta\tilde{u}^3 = 0$ is generated under the mirror transformation $u = -\tilde{u}$, which possesses the same property as the old equation (1).

The transformation (28) maps the case: $\delta > 0, \Delta \geq 0$ into the same case: $\tilde{\delta} > 0, \tilde{\Delta} \geq 0$. Therefore a new Kolmogorov-Petrovskii-Piskunov equation (29) has also five explicit exact solutions defined by

$$\tilde{u}_i(x, t) = u_i(x, t) \big|_{\mu=\tilde{\mu}, \nu=\tilde{\nu}, \delta=\tilde{\delta}}, \quad 1 \leq i \leq 5,$$

and further five new exact solutions to the old Kolmogorov-Petrovskii-Piskunov equation (1) may be presented by $\alpha\tilde{u}_i(x, t) + \beta$, $1 \leq i \leq 5$. However this transformation process hasn't given new kind of solutions to (1) beginning with the obtained five solutions, which will be shown below.

Note that we have for $\alpha > 0, \nu \geq 0$:

$$\begin{aligned} \sqrt{\tilde{\Delta}} &= \frac{1}{2}\alpha(\nu + \sqrt{\Delta}), \quad \tilde{\nu} - \sqrt{\tilde{\Delta}} = \alpha(-\nu + \sqrt{\Delta}), \quad \tilde{\nu} + \sqrt{\tilde{\Delta}} = 2\alpha\sqrt{\Delta}, & \text{when } \beta = C_+; \\ \sqrt{\tilde{\Delta}} &= \frac{1}{2}\alpha(\nu - \sqrt{\Delta}), \quad \tilde{\nu} - \sqrt{\tilde{\Delta}} = -\alpha(\nu + \sqrt{\Delta}), \quad \tilde{\nu} + \sqrt{\tilde{\Delta}} = -2\alpha\sqrt{\Delta}, & \text{when } \beta = C_-, \mu \geq 0; \\ \sqrt{\tilde{\Delta}} &= \frac{1}{2}\alpha(\sqrt{\Delta} - \nu), \quad \tilde{\nu} - \sqrt{\tilde{\Delta}} = -2\alpha\sqrt{\Delta}, \quad \tilde{\nu} + \sqrt{\tilde{\Delta}} = -\alpha(\nu + \sqrt{\Delta}), & \text{when } \beta = C_-, \mu \leq 0. \end{aligned}$$

The concrete results of the Bäcklund transformation

$$(\text{BT})_{\beta} : u(x, t) \mapsto \alpha u(x, t) \big|_{\mu=\tilde{\mu}, \nu=\tilde{\nu}, \delta=\tilde{\delta}} + \beta \quad (31)$$

may be given out for $\alpha > 0$, $\nu \geq 0$ as follows:

$$\begin{aligned}
(\text{BT})_{C_+} : \quad & u_1(x, t; \varepsilon_1, \varepsilon_4) \mapsto u_3(x, t; \varepsilon_3 = -\varepsilon_1, \varepsilon_6 = -\varepsilon_4), \\
& u_2(x, t; \varepsilon_2, \varepsilon_5) \mapsto u_2(x, t; -\varepsilon_2, -\varepsilon_5), \\
& u_3(x, t; \varepsilon_3, \varepsilon_6) \mapsto u_1(x, t; \varepsilon_1 = -\varepsilon_3, \varepsilon_4 = \varepsilon_6), \\
& u_4(x, t; A_1, A_2, A_3) \mapsto u_4(x, t; A_2, A_1, A_3), \\
& u_5(x, t; A_1, A_2, A_3) \mapsto u_5(x, t; A_3, A_2, A_1);
\end{aligned}$$

$$\begin{aligned}
(\text{BT})_{C_-, \mu \geq 0} : \quad & u_1(x, t; \varepsilon_1, \varepsilon_4) \mapsto u_2(x, t; \varepsilon_2 = -\varepsilon_1, \varepsilon_5 = \varepsilon_4), \\
& u_2(x, t; \varepsilon_2, \varepsilon_5) \mapsto u_3(x, t; \varepsilon_3 = -\varepsilon_2, \varepsilon_6 = -\varepsilon_5), \\
& u_3(x, t; \varepsilon_3, \varepsilon_6) \mapsto u_1(x, t; \varepsilon_1 = \varepsilon_3, \varepsilon_4 = -\varepsilon_6), \\
& u_4(x, t; A_1, A_2, A_3) \mapsto u_4(x, t; A_2, A_1, A_3), \\
& u_5(x, t; A_1, A_2, A_3) \mapsto u_5(x, t; A_3, A_2, A_1);
\end{aligned}$$

$$\begin{aligned}
(\text{BT})_{C_-, \mu \leq 0} : \quad & u_1(x, t; \varepsilon_1, \varepsilon_4) \mapsto u_2(x, t; \varepsilon_2 = \varepsilon_1, \varepsilon_5 = -\varepsilon_4), \\
& u_2(x, t; \varepsilon_2, \varepsilon_5) \mapsto u_1(x, t; \varepsilon_1 = \varepsilon_2, \varepsilon_4 = -\varepsilon_5), \\
& u_3(x, t; \varepsilon_3, \varepsilon_6) \mapsto u_3(x, t; -\varepsilon_3, -\varepsilon_6), \\
& u_4(x, t; A_1, A_2, A_3) \mapsto u_4(x, t; A_3, A_2, A_1), \\
& u_5(x, t; A_1, A_2, A_3) \mapsto u_5(x, t; A_2, A_1, A_3).
\end{aligned}$$

When $\beta = 0$, the transformation (28) also maps the other case: $\delta > 0$, $\Delta \leq 0$ into the same case: $\tilde{\delta} > 0$, $\tilde{\Delta} \leq 0$. We may show that $(\text{BT})_{\beta=0, \alpha>0}$ is an identity map on the set of solutions $\text{Span}\{u_i, u'_j | 1 \leq i \leq 5, j = 1, 2\}$ and that

$$\begin{aligned}
(\text{BT})_{\beta=0, \alpha<0} : \quad & u_1(x, t; \varepsilon_1, \varepsilon_4) \mapsto u_1(x, t; -\varepsilon_1, \varepsilon_4), \\
& u_2(x, t; \varepsilon_2, \varepsilon_5) \mapsto u_3(x, t; \varepsilon_3 = \varepsilon_2, \varepsilon_6 = -\varepsilon_5), \\
& u_3(x, t; \varepsilon_3, \varepsilon_6) \mapsto u_2(x, t; \varepsilon_2 = \varepsilon_3, \varepsilon_5 = -\varepsilon_6), \\
& u_4(x, t; A_1, A_2, A_3) \mapsto u_5(x, t; A_1, A_2, A_3), \\
& u_5(x, t; A_1, A_2, A_3) \mapsto u_4(x, t; A_1, A_2, A_3), \\
& u'_1(x, t; \varepsilon_1, \varepsilon_2) \mapsto u'_1(x, t; -\varepsilon_1, \varepsilon_2), \\
& u'_2(x, t; \varepsilon_1, \varepsilon_2) \mapsto u'_2(x, t; -\varepsilon_1, \varepsilon_2).
\end{aligned}$$

The rest case of $(\text{BT})_{C_{\pm}}$ may be computed similarly. It is interesting to note that $(\text{BT})_{\beta=0, \alpha<0}$ casts the solution (26) into the solution (27) and vice versa.

We point out that we may also transform a more general equation

$$w_t - w_{xx} = f(w) = a + bw + cw^2 + dw^3, \quad a \neq 0, \quad (32)$$

where a, b, c, d are real constants, into the Kolmogorov-Petrovskii-Piskunov equation (1) under the linear transformation $w = \alpha u + \beta$, $\alpha, \beta = \text{const}$. Therefore we can generate solutions to (32) by the obtained results. However we should note that the equation (32) lost the property $f(0) = 0$.

5 Discussions

The Kolmogorov-Petrovskii-Piskunov equation (1) contains the following various equations with $\mu + \nu + \delta = 0$, for which there is always the condition $\Delta = \nu^2 - 4\mu\delta = (\mu - \delta)^2 \geq 0$ and thus has five explicit solutions.

(i) The nonintegrable Newell-Whitehead [19] equation

$$u_t - u_{xx} = u - u^3 \quad (33)$$

is a special case with $\mu = -1$, $\nu = 0$, $\delta = 1$. The cases of $\mu = 0$, $\nu = -1$, $\delta = 1$ and $\mu = 1$, $\nu = -2$, $\delta = 1$ engender the equations

$$u_t - u_{xx} = u^2(1 - u), \quad (34)$$

$$u_t - u_{xx} = -u(1 - u)^2, \quad (35)$$

respectively. The above three equations are all simple generalizations of the Fisher equation. Interestingly, an another simple generalization of the Fisher equation $u_t - u_{xx} = u(1 - u)^2$ has no nonconstant solution of the form

$$u = \sum_{i=0}^M a_i \tanh^i(kx - \omega t + \xi_0), \quad u = \sum_{i=0}^M b_i \coth^i(kx - \omega t + \xi_0), \quad a_i, b_i \in R^1.$$

This is in agreement with the result in Ref. [6], where it was shown that this equation has no solution of the form $a_1 \tanh(kx - \omega t + \xi_0) + a_0$.

(ii) The case of $\mu = a$, $\nu = -(a + 1)$, $\delta = 1$ engenders the FitzHugh-Nagumo equation

$$u_t - u_{xx} = u(u - a)(1 - u), \quad (36)$$

where a is arbitrary. This equation may describe nerve pulse propagation in nerve fibers and wall motion in liquid crystals. Its solutions were discussed in [9] [12].

Our method in Section 2 is a kind of combination of the direct method [15] and the ansatz method [16] [22] and thus we term it the combined ansatz method. The idea is to make the unknown variable u to be a practicable function $g(v)$ of the ansatz unknown variable v which satisfies a differential equation solvable by quadratures. This allows us to solve a large class of physically important nonlinear equations including some nonintegrable ones, for example, $2D$ -KdV-Burgers equation [20] and 7th generalized KdV equation [17]. The crucial point is to choose the proper ansatz equations solvable by quadratures. We here list two useful ansatz equations and their solutions. The first one is the Bernoulli equation

$$v_\xi = av + bv^\alpha, \quad a, b, \alpha \in R^1, \quad ab \neq 0, \quad \alpha \neq 1. \quad (37)$$

It has a general solution

$$v = \left[-\frac{a}{b} \frac{1}{\xi_0 e^{a(1-\alpha)\xi} + 1} \right]^{\frac{1}{\alpha-1}} = \begin{cases} \left(-\frac{a}{2b}\right)^{\frac{1}{\alpha-1}}, & \text{for } \xi_0 = 0, \\ \left\{-\frac{a}{2b} \left[\tanh\left(\frac{a(\alpha-1)}{2}\xi - \frac{\ln \xi_0}{2}\right) + 1\right]\right\}^{\frac{1}{\alpha-1}}, & \text{for } \xi_0 > 0, \\ \left\{-\frac{a}{2b} \left[\coth\left(\frac{a(\alpha-1)}{2}\xi - \frac{\ln(-\xi_0)}{2}\right) + 1\right]\right\}^{\frac{1}{\alpha-1}}, & \text{for } \xi_0 < 0, \end{cases} \quad (38)$$

where ξ_0 is arbitrary. The second one is the Riccati equation

$$v_\xi = a_0 + a_1 v + a_2 v^2, \quad a_i \in R^1, \quad a_2 \neq 0. \quad (39)$$

This equation has the following solutions

$$v = -\frac{a_1}{2a_2}, \quad -\frac{1}{a_2 \xi + \xi_0} - \frac{a_1}{2a_2}, \quad (\Delta = 0); \quad (40)$$

$$v = -\frac{\varepsilon\sqrt{\Delta}}{a_2} \frac{1}{\xi_0 \exp(-\varepsilon\sqrt{\Delta}\xi) + 1} + \frac{\varepsilon\sqrt{\Delta}}{2a_2} - \frac{a_1}{2a_2} \quad (\varepsilon = \pm 1),$$

$$= \begin{cases} \frac{\varepsilon\sqrt{\Delta}}{2a_2} - \frac{a_1}{2a_2}, & \text{for } \xi_0 = 0, \\ -\frac{\sqrt{\Delta}}{2a_2} \tanh\left(\frac{\sqrt{\Delta}}{2}\xi - \frac{\varepsilon \ln \xi_0}{2}\right) - \frac{a_1}{2a_2}, & \text{for } \xi_0 > 0, \quad (\Delta > 0); \\ -\frac{\sqrt{\Delta}}{2a_2} \coth\left(\frac{\sqrt{\Delta}}{2}\xi - \frac{\varepsilon \ln(-\xi_0)}{2}\right) - \frac{a_1}{2a_2}, & \text{for } \xi_0 < 0, \end{cases} \quad (41)$$

$$v = \begin{cases} \frac{\sqrt{-\Delta}}{2a_2} \tan\left(\frac{\sqrt{-\Delta}}{2}\xi + \xi_0\right) - \frac{a_1}{2a_2}, & (\Delta < 0); \\ -\frac{\sqrt{-\Delta}}{2a_2} \cot\left(\frac{\sqrt{-\Delta}}{2}\xi + \xi_0\right) - \frac{a_1}{2a_2}, & \end{cases} \quad (42)$$

where $\Delta = a_1^2 - 4a_0a_2$ and ξ_0 is arbitrary. The Bernoulli equation and the Riccati equation have been appeared in [16] [22]. However, singular solutions of these two equations are all missing in their works.

For the general Fisher equation

$$u_t - u_{xx} + \mu u + \nu u^2 = 0, \quad \mu, \nu \in R^1, \quad (43)$$

the same combined ansätze as ones for the Kolmogorov-Petrovskii-Piskunov (1) may also result in solutions, but only a kind of solutions

$$u(x, t) = -\frac{\mu}{2\nu} - \frac{|\mu|}{4\nu} + \frac{\varepsilon_1\mu}{2\nu} v\left(\frac{\varepsilon_2\sqrt{6|\mu|}}{12}x + \frac{5\varepsilon_1\mu}{12}t\right) + \frac{|\mu|}{4\nu} v^2\left(\frac{\varepsilon_2\sqrt{6|\mu|}}{12}x + \frac{5\varepsilon_1\mu}{12}t\right), \quad (44)$$

where $v_\xi = 1 - v^2$, $\varepsilon_1, \varepsilon_2 = \pm 1$. Particularly, the Fisher elution (43) hasn't any solution of the form

$$u = \sum_{i=0}^M a_i \tan^i(kx - \omega t + \xi_0), \quad u = \sum_{i=0}^M b_i \cot^i(kx - \omega t + \xi_0), \quad a_i, b_i \in R^1. \quad (45)$$

It is worth pointing out that the Fisher equation can't be solved through the ansatz method proposed in Refs. [16] [22]. When we choose $v = \tanh(\xi + \xi_0)$ or $v = \coth(\xi + \xi_0)$, the solution (44) yields two explicit solutions, for instance, a traveling front solution

$$u(x, t) = -\frac{\mu}{2\nu} - \frac{|\mu|}{4\nu} + \frac{\varepsilon_1\mu}{2\nu} \tanh\left(\frac{\sqrt{6|\mu|}}{12}x + \frac{5\varepsilon_1\mu}{12}t + \xi_0\right) + \frac{|\mu|}{4\nu} \tanh^2\left(\frac{\sqrt{6|\mu|}}{12}x + \frac{5\varepsilon_1\mu}{12}t + \xi_0\right). \quad (46)$$

They contain two solutions of the standard Fisher equation $u_t - u_{xx} = u(1 - u)$.

Finally we remark that any Riccati equation possesses an important property: given a particular solution, its general solution may be found by quadratures. This property is named the Riccati property by Fuchssteiner and Carillo [8] and a method to construct ordinary differential equations which enjoy the Riccati property is proposed in their works [8] [5]. We may also take the differential equations proposed in [8] [5] as the basic ansatz equations. This may make more nonlinear equations to be solvable by quadratures.

Acknowledgment: One of the authors (W. X. Ma) acknowledges financial support from Alexander von Humboldt Foundation.

References

- [1] M. A. Abdelkader, Travelling wave solutions for a generalized Fisher equation, *J. Math. Anal. Appl.* **85**, 287-290 (1982).

- [2] M. J. Ablowitz and A. Zeppetella, Explicit solutions of Fisher's equation for a special wave speed, *Bull. Math. Bio.* **41**, 835-840 (1979).
- [3] D. G. Aronson and H. F. Weinberger, Nonlinear diffusion in population genetics, combustion and nerve pulse propagation, in *Partial Differential Equations and Related Topics*, edited by J. A. Goldstein, Lecture Notes in Mathematics, Vol. 446, Springer, Berlin, 5-49 (1975).
- [4] Bramson, M. D., Convergence of solutions of the Kolmogorov equation to travelling waves, *Mem. Am. Math. Soc.* **285**, 1-190 (1983).
- [5] S. Carillo and B. Fuchssteiner, Some remarks on a class of ordinary differential equations: the Riccati property, in: *Modern Group Analysis: Advance Analytical and Computational Methods in Mathematical Physics*, edited by N. H. Ibragimov et al., Kluwer, Dordrecht, 85-92 (1993).
- [6] H. Cohen, Nonlinear diffusion problems, in *Studies in Applied Mathematics*, edited by A. H. Taub, The Mathematical Association of America, 27-64 (1971).
- [7] R. A. Fisher, The wave of advance of advantageous genes, *Ann. Eugenics* **7**, 355-369 (1937).
- [8] B. Fuchssteiner and S. Carillo, A new class of nonlinear partial differential equations solvable by quadratures. in: *Analysis and Geometry*, edited by B. Fuchssteiner and W. A. J. Luxemburg, BI Wissenschaftsverlag, Mannheim-Leipzig-Wien-Zürich, 73-85 (1992).
- [9] W. Hereman, Application of a Macsyma program for the Painlevé test to the FitzHugh-Nagumo equation, in *Partially Integrable Evolution Equations in Physics*, edited by R. Conte and N. Boccara, Kluwer, Dordrecht, 585-586 (1990).
- [10] Y. Kametaka, On the nonlinear diffusion equation of Kolmogorov-Petrovskii-Piskunov type, *Osaka J. Math.* **13**, 11-66 (1976).
- [11] E. Kamke, *Differentialgleichungen: Lösungsmethoden u. Lösungen I.* Teubner, Stuttgart (1977).
- [12] T. Kawahara and M. Tanaka, Interactions of travelling fronts: an exact solution of a nonlinear diffusion equation, 311-314 *Phys. Lett. A* **97**, 311-314 (1983).
- [13] A. N. Kolmogorov, I. G. Petrovskii and N. S. Piskunov, A study of the diffusion equation with increase in the quantity of matter, and its application to a biological problem, *Bull. Moscow Univ. Math. ser. A* **1**, 1-25 (1937).
- [14] E. V. Krishnan, On some diffusion equations, *J. Phys. Soc. Jpn.* **63**, 460-465 (1994).
- [15] H. B. Lan and K. L. Wang, Exact solutions for two nonlinear equations: I, *J. Phys. A: Math. Gen.* **23**, 3923-3928 (1990).
- [16] B. Q. Lu, B. Z. Qu, Z. L. Pan and X. F. Jiang, Exact traveling wave solution of the one class of nonlinear diffusion equations, *Phys. Lett. A* **175**, 113-115 (1993).
- [17] W. X. Ma, Travelling wave solutions to a seventh order generalized KdV equation, *Phys. Lett. A* **180**, 221-224 (1993).
- [18] H. P. McKean, Application of Brownian motion to the equation of Kolmogorov-Petrovskii-Piskunov, *Comm. Pure Appl. Math.* **28**, 323-331 (1975).

- [19] A. C. Newell and J. A. Whitehead, Finite bandwidth, finite amplitude convection, *J. Fluid Mech.* **38**, 279-303 (1969).
- [20] E. J. Parkes, Exact solutions to the two-dimensional KdV-Burgers equation, *J. Phys. A: Math. Gen.* **27**, L497-L501 (1994).
- [21] X. Y. Wang, Explicit solitary wave solution to a generalized Fisher equation, *Chinese Science Bulletin* **33**, 657-659 (1988); Exact and explicit solitary wave solutions for the generalized Fisher equation, *Phys. Lett. A* **131**, 277-279 (1988).
- [22] Z. J. Yang, Traveling wave solutions to nonlinear evolution and wave equations, *J. Phys. A: Math. Gen.* **27**, 2837-2855 (1994).